



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Also

$$\begin{aligned} d\lambda_i &= \frac{h_i dt}{r_i^2} = h_i \frac{\rho^2}{r_i^2} d\psi \\ &= \frac{h_i a_i d\psi}{(\psi + a_i)^2 + h_i^2}, \\ \tan(\lambda_i + \beta_i) &= \frac{a_i}{h_i}(\psi + a_i). \end{aligned}$$

In the second place, let  $A^2$  be negative. Here it is only necessary in some places to accomplish the integrations by the aid of hyperbolic cosines instead of circular.

The differential equations of this problem, in the case where the *radii* are supposed to describe no areas, were first integrated by Binet.\* But the addition, to the forces, of the terms arising from centrifugal action, much enhances the interest of the problem.

\*See Liouville, *Journal de Mathématiques*, First Ser. Tome II. p. 457.



## ON THE FOCAL CHORD OF A PARABOLA.

By PROF. R. H. GRAVES, Chapel Hill, N. C.

Let  $y^2 = 4ax$  be the equation to a parabola,  $s$  its focus, and  $PSP'$  a focal chord. Let the tangent and normal at  $P'$  meet the diameter through  $P$  at  $M$  and  $N$ .

It may be easily proved that  $PM = PN = PP'$  and that a similar property holds for the tangent and normal at  $P$ .

Therefore, if two equal rhombs be constructed on  $PP'$  having two other sides of each parallel to the axis, their diagonals are tangents and normals at  $P$  and  $P'$ ; and the tangent at one point is parallel to the normal at the other.

Each normal chord divides the other in the ratio 1 : 3.

The chord joining the other ends of the normal chords is parallel to  $PP'$  and three times as long.

A line perpendicular to  $PP'$  at  $S$ , and terminated by this parallel chord and the pole of  $PP'$ , is divided by  $S$  in the ratio 1 : 4.

Hence the locus of the foot of the perpendicular dropped from  $S$  on the parallel chord is a right line, whose equation is  $x = 9a$ .

Hence the envelope of the parallel chord is a confocal parabola, having for its equation  $y^2 = 32a(9a - x)$ .

It cuts the original parabola orthogonally where it is cut by its evolute.